



# How Vascular Density Correlates to Visual Field Defects in Posner Schlosman Syndrome - A Case Study

**Bui Thi Van Anh\***; **Duong Phuong Loan**; **Bui Viet Hung**

High-tech Eye Center, Hanoi Tam Anh General Hospital, Hanoi, Vietnam.

## \*Corresponding Authors: Bui Thi Van Anh

High-tech Eye Center, Hanoi Tam Anh General Hospital,  
Hanoi, Vietnam.

Email: buivananh@yahoo.com &  
anhbtv1@tamanhospital.vn

## Abstract

During high IOP term, vessel density diffusely reduces, particularly on the superficial layer, but doesn't correspond to visual field defect. On the other hand, during the controlled IOP term, the loss of vessel area of ONH well corresponds to the VF defect.

Swept-source OCTA enables the acquisition of images down to the lamina cribrosa, as well as the clear separation of the vascular distribution between the layers. Therefore, SS-OCTA imaging can be used to determine the location of changes in vascular density, thereby identifying the source of damaged vessels or the source of pathology.

Received: Jan 02, 2026

Accepted: Feb 02, 2026

Published Online: Feb 09, 2026

Journal: Annals of Ophthalmology and Visual Sciences

Publisher: MedDocs Publishers LLC

Online edition: <http://meddocsonline.org/>

Copyright: © Van Anh BT (2026). *This Article is distributed under the terms of Creative Commons Attribution 4.0 International License*

**Keywords:** ONH; SS-OCTA; DRI Triton; Vessel density; Superficial layer; Deeper layer.

## Introduction

The optic nerve head and retina are supplied by two vascular systems: the uveal vessels and the central retina vessels, which originate from the ophthalmic artery, a branch of the internal carotid artery.

The circulation in the Optic Nerve Head (ONH) is divided into superficial layers and deeper layers (prelaminar, lamina cribrosa, post-lamina area, and outer retinal layers). The blood of the superficial layer, supplied via small branches of the central retinal artery by creating a radial capillary network surrounding the optic disc, nurtures the nerve fiber layer and the inner retina (from ILM to IPL/ILN layer). The prelaminar area, lamina cribrosa, and outer retinal layers (from OS/RPE to Bruch membrane), however, are directly supplied by the capillaries that are exceptionally long and originate from short posterior ciliary arteries or branches from the choroidal arteries. Unlike the arte-


rial supply, venous drainage of the ONH is through the central retinal vein.

DRI Triton SS-OCT device (Topcon, Tokyo, Japan) can analyze Vessel Density (VD) of the Optic nerve head area as deep as the lamina cribrosa, something that SDOCT is hard to do. By applying the Macular Angio protocol for scanning the vessels at ONH, we can get the vessel's information inside a 1mm circle from the central retinal vessels, which we cannot reach by using only the Disc Angio protocol. In clinical practice, we use the Disc Angio protocol to survey the *peripapillary vessel* located between two concentric circles: a 2mm circle and a 4mm circle, with the center coinciding with the center of the optic disc. The Macular Angio protocol is used to get information about the *Central ONH vessel* within the optic disc margin or a 1 mm in diameter center circle. The Macular Angio protocol is also used to analyze the juxta-papillary vessels, which are located immediately adjacent to the optic disc margin, 1 mm wide, measured outwards from



**Cite this article:** Van Anh BT, Loan DP, Hung BV. How Vascular Density Correlates to Visual Field Defects in Posner Schlosman Syndrome-A Case Study. *Ann Ophthalmol Vis Sci.* 2026; 9(1): 1053.

the central optic disc survey area (a circle with a diameter of 3 mm centered on the optic disc margin).



**Figure 1:** Vascular System of Optic Nerve Head.

Optic nerve head vessel density of a normal person (19-year-old male, left eye)

Photos were taken using the Disc Angio protocol: 1a. vessel density of whole retinal thickness; 1b. vessel density of the superficial part of retina (from ILM to IPL/INL); 1c. vessel density of the deeper part of the retina (from OS/RPE to Bruch membrane)

Photos were taken using the Macular Angio protocol: 1d. vessel density of whole retinal thickness; 1e. vessel density of the superficial part of retina (from ILM to IPL/INL); 1f. vessel density of the deeper part of the retina (from OS/RPE to Bruch membrane)


**Clinical Case**

Had finished the treatment for shingles on the right half of the face 10 days ago, a 56-year-old female came to check her eyes because of pain in the right eye, lasting for 3 days

Her acuity vision changed rapidly to 20/100 with an IOP was 60 mmHg on the right eye, while the left eye was almost normal at 20/25 and IOP was 11 mmHg. There were some mutton precipitates and edema cornea on the right. Fundus, RNFL on OCT, and visual field were almost normal except for a small nasal step lesion on the right VF (Figure 2).



**Figure 2:** OCT and VF of right eye.




**Figure 3:** Vessel density of ONH in the right eye: diffuse ischemia in the nasal half of the superficial vascular layer (B-E), the deep blood vessels were almost unaffected (C-F).

OCTA images were taken to analyze the vascular system of ONH. The vascular density distribution plan showed a diffuse ischemia in the nasal half of the right eye, completely inconsistent with the damage in the visual field. This condition only occurred in the superficial vascular layer; the deeper vessels were almost unaffected.

The patient was diagnosed with Posner-Schlossman syndrome in the right eye. Having been treated with steroids and Azarga, and Acetazolamide 0.25g 2 tablets/day, the right eye's IOP decreased to 11 mmHg.

After 3 days, the superficial optic nerve ischemia at the optic disc and around the optic disc significantly improved beyond the ischemic band at 5 o'clock. The location of this fixed ischemic lesion corresponded to the visual field defect of the right eye.



**Figure 4:** Distribution of ONH's vessel in the superficial layer when IOP 60 mmHg (A,B) and IOP 11 mmHg (C,D).

**Discussion**

Retinal blood flow depends not only on the perfusion pressure, which is the difference between retinal arterial and retinal venous pressure, but also on the level of intraocular pressure. The local resistance around ONH was high when IOP was 60mmHg and was reduced when IOP was 11 mmHg, resulting in a reverse increase in retinal vascular perfusion. During the high IOP term, some vessels were actually occluded and manifested as an avascular zone on the vascular distribution image when IOP was normal (Figure 4), which corresponded to VF defects (Figure 3). We hypothesized that, in the early stage of glauco-

ma, only the vessels on the superficial layer are damaged, but in the end stage, both vessels on the superficial and deep layers are damaged. Only the non-perfusion zone in the normal IOP term corresponds with defects of VF.

#### Author declarations

#### Acknowledge

Thank you for the participation of my colleagues, Optometrist. Duong Phuong Loan; Dr. Nguyen Duc Doanh, Dr. Bui Viet Hung (High-tech Eye Center, Hanoi Tam Anh General Hospital).

#### References

1. Hofman P, Hoyng P, VanderWerf F, Vrensen GF, Schlingemann RO. Lack of blood–brain barrier properties in microvessels of the prelaminar optic nerve head. *Invest Ophthalmol Vis Sci.* 2001; 42: 895–901.
2. Flammer J, Konieczka K, Flammer AJ. The primary vascular dysregulation syndrome: implications for eye diseases. *EPMA J.* 2013; 4: 14.
3. Sossi N, Anderson DR. Blockage of axonal transport in optic nerve induced by elevation of intraocular pressure: effect of arterial hypertension induced by angiotensin I. *Arch Ophthalmol.* 1983; 101: 94–97.
4. Flammer J, Orgül S, Costa VP, et al. The impact of ocular blood flow in glaucoma. *Prog Retin Eye Res.* 2002; 21: 359–93.
5. Bojikian KD, Chen CL, Wen JC, Zhang Q, Xin C, Gupta D, et al. Optic disc perfusion in primary open-angle and normal tension glaucoma eyes using optical coherence tomography-based microangiography. *PLoS One.* 2016; 11: e0154691.
6. Lee EJ, Lee KM, Lee SH, Kim TW. OCT angiography of the peripapillary retina in primary open-angle glaucoma. *Invest Ophthalmol Vis Sci.* 2016; 57: 6265–70.
7. Chen CL, Zhang A, Bojikian KD, et al. Peripapillary retinal nerve fiber layer vascular microcirculation in glaucoma using optical coherence tomography-based microangiography. *Invest Ophthalmol Vis Sci.* 2016; 57: 475–85.
8. Campbell JP, Zhang M, Hwang TS, et al. Detailed vascular anatomy of the human retina by projection-resolved optical coherence tomography angiography. *Sci Rep.* 2017; 7: 42201.
9. Yarmohammadi A, Zangwill LM, Diniz-Filho A, et al. Relationship between optical coherence tomography angiography vessel density and severity of visual field loss in glaucoma. *Ophthalmology.* 2016; 123: 2498–508.
10. Spaide RF, Fujimoto JG, Waheed NK, Sadda SR, Staurengi G. Optical coherence tomography angiography. *Prog Retin Eye Res.* 2018; 64: 1–55.