



# Disseminated Blastomycosis

Michael Stewart DO<sup>1\*</sup>; Rakin Solaiman MD<sup>1</sup>

Internal Medicine Residency Spokane, Spokane, Washington.

**\*Corresponding Author(s): Michael Stewart**

Internal Medicine Residency Spokane, Spokane, Washington.

Email: Michael.stewart2@providence.org

Received: Mar 10, 2024

Accepted: Mar 26, 2024

Published Online: Mar 29, 2024

Journal: Journal of Case Reports and Medical Images

Publisher: MedDocs Publishers LLC

Online edition: <http://meddocsonline.org/>

Copyright: © Stewart M (2024). This Article is distributed under the terms of Creative Commons Attribution 4.0 International License

## Clinical Image Description

A 24-year-old male presented with a 2-year history of an insidiously expanding lesion to his left lateral thigh (Panel A) and widespread pink annular scaly plaques with central hypopigmentation to torso and extremities. He had not been seen for the lesion or rash until being hospitalized for a new cough and painful erythematous joint of right knee. Arthrocentesis showed 21,126 nucleated cells/uL with 98% neutrophils, and MRI demonstrated a 1.5x2.4x6.7cm right femoral bone abscess with concomitant large joint effusion and enhancing synovitis (Panel B). Orthopedics was consulted with subsequent right knee arthrotomy, synovectomy, and partial excision and debridement of right distal femur. Two separate lesion punch biopsies, joint effusion analysis, and bone abscess evaluation all demonstrated budding yeast (Panel C).

Infectious disease started empiric voriconazole. When cultures were confirmed as Blastomycosis dermatitidis, treatment was changed to itraconazole. At 4-week follow-up, there was healthy remodeling and wound contracture of thigh lesion.

